

Reray ... Narayanan R.

Sharma, H.W.  
Anticancer Research

1996 16 61  
RC 201.A/1868

Tam J. Immunol. 1997

158 200-208

OR 180.46

Blackburn G.H. (1990)

JBC 265:5919

Ggunter

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:sssptal635jxs

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

\* \* \* \* \* Welcome to STN International

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 Jan 25 BLAST(R) searching in REGISTRY available in STN on the Web  
NEWS 3 Jan 29 FSTA has been reloaded and moves to weekly updates  
NEWS 4 Feb 01 DKILIT now produced by FIZ Karlsruhe and has a new update frequency  
NEWS 5 Feb 19 Access via Tymnet and SprintNet Eliminated Effective 3/31/02  
NEWS 6 Mar 08 Gene Names now available in BIOSIS  
NEWS 7 Mar 22 TOXLIT no longer available  
NEWS 8 Mar 22 TRCTHERMO no longer available  
NEWS 9 Mar 28 US Provisional Priorities searched with P in CA/CAPLUS and USPATFULL  
NEWS 10 Mar 28 LIPINSKI/CALC added for property searching in REGISTRY  
NEWS 11 Apr 02 PAPERCHEM no longer available on STN. Use PAPERCHEM2 instead.  
NEWS 12 Apr 08 "Ask CAS" for self-help around the clock  
NEWS 13 Apr 09 BEILSTEIN: Reload and Implementation of a New Subject Area  
NEWS 14 Apr 09 ZDB will be removed from STN  
NEWS 15 Apr 19 US Patent Applications available in IFICDB, IFIPAT, and IFIUDB  
NEWS 16 Apr 22 Records from IP.com available in CAPLUS, HCAPLUS, and ZCAPLUS  
NEWS 17 Apr 22 BIOSIS Gene Names now available in TOXCENTER  
NEWS 18 Apr 22 Federal Research in Progress (FEDRIP) now available  
NEWS 19 May 31 PCTFULL to be reloaded. File temporarily unavailable.  
NEWS 20 Jun 03 New e-mail delivery for search results now available  
  
NEWS EXPRESS February 1 CURRENT WINDOWS VERSION IS V6.0d,  
CURRENT MACINTOSH VERSION IS V6.0a(ENG) AND V6.0Ja(JP),  
AND CURRENT DISCOVER FILE IS DATED 05 FEBRUARY 2002  
  
NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\* \* \* \* \* STN Columbus \* \* \* \* \*

FILE 'HOME' ENTERED AT 16:03:54 ON 07 JUN 2002

=> FIL BIOSIS MEDLINE SCISEARCH CA  
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'BIOSIS' ENTERED AT 16:04:08 ON 07 JUN 2002  
COPYRIGHT (C) 2002 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'MEDLINE' ENTERED AT 16:04:08 ON 07 JUN 2002

FILE 'SCISEARCH' ENTERED AT 16:04:08 ON 07 JUN 2002  
COPYRIGHT (C) 2002 Institute for Scientific Information (ISI) (R)

FILE 'CA' ENTERED AT 16:04:08 ON 07 JUN 2002  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

<-----User Break----->

u  
SEARCH ENDED BY USER

=> s SP1 or (NF Kappa B) or EGR1 or AP2  
L1 65870 SP1 OR (NF KAPPA B) OR EGR1 OR AP2

=> s l1 and aptamer  
L2 21 L1 AND APTAMER

=> s l2 and g-rich aptamer  
L3 0 L2 AND G-RICH APTAMER

=> s l2 and g-rich  
L4 1 L2 AND G-RICH

=> d l4 ti

L4 ANSWER 1 OF 1 CA COPYRIGHT 2002 ACS  
TI **G-rich** oligonucleotides binding transcription factors  
involved in inflammatory responses for the treatment of inflammatory  
disease

=> d l4 bib

L4 ANSWER 1 OF 1 CA COPYRIGHT 2002 ACS  
AN 129:117842 CA  
TI **G-rich** oligonucleotides binding transcription factors  
involved in inflammatory responses for the treatment of inflammatory  
disease  
IN Tam, Robert  
PA ICN, Pharmaceuticals, Inc., USA  
SO PCT Int. Appl., 43 pp.  
CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	----	-----	-----
PI WO 9829430	A1	19980709	WO 1997-US23927	19971219
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US,			

UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM  
 RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI,  
 FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM,  
 GA, GN, ML, MR, NE, SN, TD, TG  
 AU 9857200 A1 19980731 AU 1998-57200 19971219  
 EP 968226 A1 20000105 EP 1997-953460 19971219  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, FI  
 CN 1242775 A 20000126 CN 1997-181056 19971219  
 BR 9714438 A 20000321 BR 1997-14438 19971219  
 JP 2002512599 T2 20020423 JP 1998-530233 19971219  
 NO 9903170 A 19990825 NO 1999-3170 19990625  
 PRAI US 1996-34509P P 19961227  
 WO 1997-US23927 W 19971219

=> d his

(FILE 'HOME' ENTERED AT 16:03:54 ON 07 JUN 2002)

FILE 'BIOSIS, MEDLINE, SCISEARCH, CA' ENTERED AT 16:04:08 ON 07 JUN 2002  
 L1 65870 S SP1 OR (NF KAPPA B) OR EGR1 OR AP2  
 L2 21 S L1 AND APTAMER  
 L3 0 S L2 AND G-RICH APTAMER  
 L4 1 S L2 AND G-RICH

=> dup l2

ENTER REMOVE, IDENTIFY, ONLY, OR (?):remove

PROCESSING COMPLETED FOR L2

L5 11 DUP REMOVE L2 (10 DUPLICATES REMOVED)

=> d l5 bib abs 1-11

L5 ANSWER 1 OF 11 CA COPYRIGHT 2002 ACS  
 AN 135:285337 CA  
 TI Methods for identifying peptide **aptamers** capable of altering a  
 cell phenotype  
 IN Benson, John D.; Vincent, Sylvie Magali; Brasher, Bradley Bryan  
 PA Enanta Pharmaceuticals, Inc., USA  
 SO PCT Int. Appl., 41 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001075178	A2	20011011	WO 2001-US10953	20010404
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
PRAI US 2000-194722P	P	20000404		
AB The invention provides methods and compns. for screening and identifying peptide <b>aptamers</b> that can modulate a cell phenotype and further, can be used for the treatment of a disease involving a misregulated cell phenotype, such as, for example, a cancer. The method involves contacting cells with a library of expressible nucleic acid sequences encoding random peptide <b>aptamers</b> , selecting at least one cell having an altered				

phenotype, and identifying the expressed **aptamers**.

L5 ANSWER 2 OF 11 MEDLINE DUPLICATE 1  
AN 2001301737 MEDLINE  
DN 21226230 PubMed ID: 11327864  
TI In vivo recognition of an RNA **aptamer** by its transcription factor target.  
AU Cassiday L A; Maher L J 3rd  
CS Department of Biochemistry and Molecular Biology, Mayo Foundation, Rochester, Minnesota 55905, USA.  
NC GM54411 (NIGMS)  
SO BIOCHEMISTRY, (2001 Feb 27) 40 (8) 2433-8, 38 3168  
Journal code: A0G; 0370623. ISSN: 0006-2960.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200105  
ED Entered STN: 20010604  
Last Updated on STN: 20010604  
Entered Medline: 20010531  
AB In vitro-selected RNA **aptamers** are potential inhibitors of disease-related macromolecules. Our laboratory previously isolated an RNA **aptamer** that specifically binds to the human transcription factor NF-kappaB. This RNA **aptamer** competitively inhibits DNA binding by NF-kappaB in vitro. In the study presented here, this **aptamer** was tested for binding to the p50 homodimer form of NF-kappaB (p50(2)) in eukaryotic cells using a yeast three-hybrid system. We show that the alpha-p50 RNA **aptamer** selectively binds recombinant p50(2) expressed in yeast, demonstrating in vivo recognition of an in vitro-selected RNA **aptamer** by its protein target. This result suggests that RNA decoys might be used to inhibit the function of DNA-binding proteins in vivo.

L5 ANSWER 3 OF 11 SCISEARCH COPYRIGHT 2002 ISI (R)  
AN 2001:989140 SCISEARCH  
GA The Genuine Article (R) Number: 500KC  
TI Intramers as promising new tools in functional proteomics  
AU Famulok M (Reprint); Blind M; Mayer G  
CS Univ Bonn, Kekule Inst Organ Chem & Biochem, Gerhard Domagk Str 1, D-53121 Bonn, Germany (Reprint); Univ Bonn, Kekule Inst Organ Chem & Biochem, D-53121 Bonn, Germany; NascaCell GmbH, D-82327 Tutzing, Germany  
CYA Germany  
SO CHEMISTRY & BIOLOGY, (OCT 2001) Vol. 8, No. 10, pp. 931-939.  
Publisher: CURRENT BIOLOGY LTD, 84 THEOBALDS RD, LONDON WC1X 8RR, ENGLAND. ISSN: 1074-5521.  
DT General Review; Journal  
LA English  
REC Reference Count: 67  
\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*  
AB **Aptamers** are valuable tools for studying numerous aspects of biological processes, opening up new experimental opportunities to analyse the function of a wide range of cellular molecules. Functional RNA molecules can be rapidly selected in vitro from complex combinatorial mixtures of different sequences. Recently, it was shown that in vitro selection processes can be automated: the First generation selection robots will soon mean **aptamers** for several targets can be isolated in parallel within days rather than weeks. **Aptamers** not only exhibit highly specific molecular recognition properties but are also able to modulate the function of their cognate targets in a highly specific manner by agonistic or antagonistic mechanisms. These properties prompted the development of novel technologies to exploit the use of **aptamers** to modulate distinct functions of biological targets.

Recent controlled expression of **aptamers** inside cells demonstrated their impressive potential as rapidly generated intracellular inhibitors of biomolecules. Intracellularly applied **aptamers** are also called 'intramers'. Here we discuss recent developments and strategies for intramer-based technologies that have the potential to greatly facilitate characterisation of unknown protein functions in the context of their natural expression status in vivo. Thus, intramer-based technologies offer many promising applications in functional genomics, proteomics and drug discovery. (C) 2001 Elsevier Science Ltd. All rights reserved.

L5 ANSWER 4 OF 11 CA COPYRIGHT 2002 ACS  
 AN 132:322077 CA  
 TI prepn. of **aptamers** contg. thymidine phosphorodithioates and their binding to nuclear factor- $\kappa$ B  
 IN Gorenstein, David G.; Aronson, Judy; Luxon, Bruce; Herzog, Norbert  
 PA Board of Regents the University of Texas System, USA  
 SO PCT Int. Appl., 63 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000024404	A1	20000504	WO 1999-US24058	19991026
	W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
	EP 1143980	A1	20011017	EP 1999-956560	19991026
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
PRAI	US 1998-105600P	P	19981026		
	WO 1999-US24058	W	19991026		

AB The present invention provides a method for concurrent achiral nucleotide modification and amplification using PCR. Provided by this method are NF- $\kappa$ B specific thioaptamers of novel sequence. This invention further provides methods of post-selection **aptamer** modification wherein one or more selected nucleotides of **aptamers** of known sequence are substituted with modified achiral nucleotides, particularly achiral thiophosphate nucleotides, wherein the substitution results in increased nuclease resistance while retaining binding efficiency and selectivity. Thio-substitution of post-selection **aptamers** with specificity for the nuclear factor, **NF- $\kappa$ B**, produced in accordance with this method have increased binding affinity and specificity in addn. to nuclease resistance. Also provided are methods for fractionating oligonucleotides depending on their degree of thio-substitution by anion exchange chromatog.

RE.CNT 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 5 OF 11 CA COPYRIGHT 2002 ACS  
 AN 134:189664 CA  
 TI DNA binding properties of the Arabidopsis floral development protein AINTEGUMENTA  
 AU Nole-Wilson, Staci; Krizek, Beth A.  
 CS Department of Biological Sciences, University of South Carolina, Columbia, SC, 29208, USA  
 SO Nucleic Acids Research (2000), 28(21), 4076-4082

CODEN: NARHAD; ISSN: 0305-1048

PB Oxford University Press

DT Journal

LA English

AB The Arabidopsis protein AINTEGUMENTA (ANT) is a member of a plant-specific family of transcription factors (**AP2**/EREBP) that share either one or two copies of an approx. 70 amino acid region called the **AP2** repeat. DNA binding activity has been demonstrated previously for members of this family contg. a single **AP2** repeat. Using an in vitro selection procedure, the DNA binding specificity of the two **AP2** repeat contg. protein ANT was found to be 5'-gCAC(A/G)N(A/T)Tccc(a/g)ANG(c/t)-3'. This consensus site is much longer than sites recognized by proteins contg. a single **AP2** repeat and neither **AP2** repeat of ANT was alone capable of binding to the selected sequences, suggesting that both **AP2** repeats make DNA contacts. ANT binds to these DNA sequences as a monomer but a higher order complex is also obsd. at high protein concns. The ANT consensus site shows some similarity to the C-repeat/DRE elements bound by proteins that contain a single **AP2** repeat, and we find that ANT binds weakly to such sites. We propose a model in which each **AP2** repeat of ANT contacts adjacent sites within the consensus sequence. Our results suggest that the **AP2** repeat can be utilized in different ways for DNA binding.

RE.CNT 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5 ANSWER 6 OF 11 MEDLINE DUPLICATE 2  
AN 2000077846 MEDLINE  
DN 20077846 PubMed ID: 10612599  
TI **Aptamers** containing thymidine 3'-O-phosphorodithioates:  
synthesis and binding to nuclear factor-kappaB.  
AU Yang X; Fennelwald S; Luxon B A; Aronson J; Herzog N K; Gorenstein D G  
GS Sealy Center for Structural Biology, Department of Human Biological  
Chemistry & Genetics, The University of Texas Medical Branch at Galveston,  
77555-1157, USA.  
NC 1C06CA59098 (NCI)  
AI27744 (NIAID)  
SO BIOORGANIC AND MEDICINAL CHEMISTRY LETTERS, (1999 Dec 6) 9 (23) 3357-62.  
Journal code: C8B; 9107377. ISSN: 0960-894X. QP501.37  
CY ENGLAND: United Kingdom  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200001  
ED Entered STN: 20000124  
Last Updated on STN: 20000124  
Entered Medline: 20000113  
AB **Aptamers** targeting NF-kappaB containing thymidine  
3'-O-phosphorodithioates in selected positions of an oligonucleotide  
duplex were synthesized. Binding affinities to NF-kappaB varied with the  
number and positions of the dithioate backbone substitutions. One of the  
**aptamers** showed specific binding to a single NF-kappaB dimer in  
cell culture extracts.

L5 ANSWER 7 OF 11 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.DUPLICATE  
3  
AN 1999:174859 BIOSIS  
DN PREV199900174859  
TI Selection and characterization of an RNA decoy for transcription factor  
NF-kappaB.  
AU Lebruska, Lori L.; Maher, L. James, III (1)  
CS (1) Department of Biochemistry and Molecular Biology, Mayo Foundation, 200  
First St., S.W., Guggenheim 16, Rochester, MN, 55905 USA

SO Biochemistry, (March 9, 1999) Vol. 38, No. 10, pp. 3168-3174.  
ISSN: 0006-2960.

DT Article

LA English

AB Despite their chemical similarity, DNA and RNA sequences typically adopt very different structures within cells and are recognized by different proteins. However, a few interesting examples of proteins with dual specificity for DNA and RNA have previously been noted. These observations raise the possibility that RNA surrogates might be identified for many transcription factors that normally bind DNA. As an initial test of this novel concept, we used in vitro selection to isolate a small RNA **aptamer** that binds with nanomolar affinity to human transcription factor NF-kappaB, a key regulator of inflammation, HIV-1 gene expression, and apoptosis. Selected RNAs contain a 31-nucleotide core domain that was shown by mutation and deletion analyses to be necessary and sufficient for NF-kappaB binding. Neither DNA nor 2'-O-methyl RNA analogues of the **aptamer** bound NF-kappaB. The results of competition experiments demonstrate that binding of the RNA **aptamer** blocks the ability of NF-kappaB to bind duplex DNA. Expression of this **aptamer** structure within heterologous nuclear RNA transcripts may provide a new strategy to inhibit NF-kappaB function in vivo. **Aptamers** that inhibit transcription factors might be useful in a variety of applications.

L5 ANSWER 8 OF 11 SCISEARCH COPYRIGHT 2002 ISI (R)

AN 1999:672823 SCISEARCH

GA The Genuine Article (R) Number: 230HU

TI Antisense strategies: functions and applications in immunology

AU Varga L V; Toth S; Novak I; Falus A (Reprint)

CS SEMMELWEIS UNIV MED, DEPT GENET CELL & IMMUNOBIOLOG, H-1085 BUDAPEST, HUNGARY (Reprint); SEMMELWEIS UNIV MED, DEPT GENET CELL & IMMUNOBIOLOG, H-1085 BUDAPEST, HUNGARY

CYA HUNGARY

SO IMMUNOLOGY LETTERS, (3 AUG 1999) Vol. 69, No. 2, pp. 217-224.  
Publisher: ELSEVIER SCIENCE BV, PO BOX 211, 1000 AE AMSTERDAM, NETHERLANDS.  
ISSN: 0165-2478.

DT General Review; Journal

FS LIFE

LA English

REC Reference Count: 43

\*ABSTRACT IS AVAILABLE IN THE ALL AND IALL FORMATS\*

AB The ability to manipulate gene expression by means of exogenously administered oligodeoxynucleotides complementary to specific sequences in the genome is clearly going to impact on many fields of biology and medicine including basic and clinical immunology. Also endogenously generated antisense RNA species are important in regulating gene expression. Antisense RNA has become a widely used tool for analysis of gene function and holds great promise for therapeutic use in the future. Thus, inhibition can take place on different levels (transcription, translation and **aptamer** binding). Avoiding unspecific reactions one has to use controls and well-designed oligonucleotides. Based on the studies described in this review, antisense oligonucleotides hold a great promise as a novel class of therapeutic agents in immunology as well as in oncology, neurology and viral infections. (C) 1999 Elsevier Science B.V. All rights reserved.

L5 ANSWER 9 OF 11 CA COPYRIGHT 2002 ACS

AN 129:117842 CA

TI G-rich oligonucleotides binding transcription factors involved in inflammatory responses for the treatment of inflammatory disease

IN Tam, Robert

PA ICN, Pharmaceuticals, Inc., USA

SO PCT Int. Appl., 43 pp.  
 CODEN: PIXXD2  
 DT Patent  
 LA English  
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 9829430	A1	19980709	WO 1997-US23927	19971219
	W:				
	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW:				
	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
	AU 9857200	A1	19980731	AU 1998-57200	19971219
	EP 968226	A1	20000105	EP 1997-953460	19971219
	R:				
	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
	CN 1242775	A	20000126	CN 1997-181056	19971219
	BR 9714438	A	20000321	BR 1997-14438	19971219
	JP 2002512599	T2	20020423	JP 1998-530233	19971219
	NO 9903170	A	19990825	NO 1999-3170	19990625
PRAI	US 1996-34509P	P	19961227		
	WO 1997-US23927	W	19971219		

AB Oligonucleotides that specifically bind to the DNA binding site of proteins such as **Sp1** and **Sp1**-related proteins involved in the regulation of expression of genes for costimulatory mols. such as CD28 and cytokines such as IL-2 and GMCSF are described. The oligonucleotides have at least two G-rich sequences of 3-4 bases sepd. by 3-6 nucleotides. These oligonucleotides compete with the endogenous sites binding these regulatory proteins of genes for involved in the regulation of T-cell activation. This serves to modulate gene expression by preventing transcription of the gene. **Aptamers** are administered to provide therapies for diseases which involve aberrant T-cell activation such as psoriasis, Type I (insulin-dependent) diabetes mellitus, multiple sclerosis, autoimmune uveitis, rheumatoid arthritis, systemic lupus erythematosus, inflammatory bowel disease (Crohn's and ulcerative colitis), and septic shock and to regulate normal T-cell activation such as in allograft rejection.

L5 ANSWER 10 OF 11 SCISEARCH COPYRIGHT 2002 ISI (R)  
 AN 1998:235158 SCISEARCH  
 GA The Genuine Article (R) Number: ZA970  
 TI Biochemistry and molecular genetics 1997  
 AU Buchner J (Reprint); Seckler R  
 SO NACHRICHTEN AUS CHEMIE TECHNIK UND LABORATORIUM, (FEB 1998) Vol. 46, No. 2, pp. 182-195.  
 Publisher: WILEY-V C H VERLAG GMBH, POSTFACH 10 11 61, D-69451 WEINHEIM, GERMANY.  
 ISSN: 0341-5163.  
 DT General Review; Journal  
 LA German  
 REC Reference Count: 214

L5 ANSWER 11 OF 11 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.DUPLICATE  
 4  
 AN 1996:456863 BIOSIS  
 DN PREV199699179219  
 TI Dissecting protein:protein interactions between transcription factors with an RNA **aptamer**.  
 AU Tian, Yu; Adya, Neeraj; Wagner, Susanne; Giam, Chou-Zen; Green, Michael



R.; Ellington, Andrew D. (1)  
CS (1) Dep. Chem., Indiana Univ., Bloomington, IN 47405 USA  
SO RNA (New York), (1995) Vol. 1, No. 3, pp. 317-326.  
ISSN: 1355-8382. *QP623.R52*

DT Article

LA English

AB Nucleic acid **aptamers** isolated from random sequence pools have generally proven useful at inhibiting the interactions of nucleic acid binding proteins with their cognate nucleic acids. In order to develop reagents that could also be used to study protein:protein interactions, we have used in vitro selection to search for RNA **aptamers** that could interact with the transactivating protein Tax from human T-cell leukemia virus. Tax does not normally bind to nucleic acids, but instead stimulates transcription by interacting with a variety of cellular transcription factors, including the cyclic AMP-response element binding protein (CREB), **NF-kappa-B**, and the serum response factor (SRF). Starting from a pool of greater than 10<sup>13</sup> different RNAs with a core of 120 random sequence positions, RNAs were selected for their ability to be co-retained on nitrocellulose filters with Tax. After five cycles of selection and amplification, a single nucleic acid species remained. This **aptamer** was found to bind Tax with high affinity and specificity, and could disrupt complex formation between Tax and CREB. Further assays revealed that the RNA could also disrupt complex formation between Tax and **NF-kappa-B**, but not with SRF. The differential effects of our **aptamer** probe on protein:protein interactions suggest a model for how the transcription factor binding sites on the surface of the Tax protein are organized. This model is consistent with data from a variety of other studies.

=>

---Logging off of STN---

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	50.89	51.10
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-2.36	-2.36

STN INTERNATIONAL LOGOFF AT 16:08:07 ON 07 JUN 2002